

Quenched QCD with domain wall fermions

P. Chen, N. Christ, G. Fleming, A. Kaehler, C. Malureanu, R. Mawhinney*, G. Siegert†, C. Sui, P. Vranas‡, and Y. Zhestkov^a

^aDepartment of Physics, Columbia University, New York, NY 10027, USA

We report on simulations of quenched QCD using domain wall fermions, where we focus on basic questions about the formalism and its ability to produce expected low energy hadronic physics for light quarks. The work reported here is on quenched $8^3 \times 32$ lattices at $\beta = 5.7$ and 5.85 , using values for the length of the fifth dimension between 10 and 48. We report results for parameter choices which lead to the desired number of flavors, a study of undamped modes in the extra dimension and hadron masses.

1. INTRODUCTION

Domain wall fermions [1] were originally proposed as a technique for putting chiral fermions on the lattice. For fermions coupled to a vector gauge theory (like QCD), this approach has given a way to treat massless (or at least controllably small mass) fermions, while preserving the full continuum symmetry group. Subsequent improvements to the original formulation [2] made simulations more practical and studies of the Schwinger model [3], including dynamical fermion effects, and quenched QCD [4] have been very encouraging. Here we report a reasonably systematic study of low energy, non-anomalous QCD physics using domain wall fermions. For a review and further references, see [5].

2. BASIC QUESTIONS

For quenched domain wall simulations, there are three input parameters for the fermions: m_f , the explicit four-dimensional bare quark mass; L_s , the extent of the lattice in the fifth dimension and m_0 , the five-dimensional bare quark mass. Except for different symbols for the three parameters listed above, our conventions follow [2].

A first basic question involves choosing parameters so that the desired number of flavors of light fermions is being studied. For free fermions, one easily sees that for $m_0 < 0$ there are no light surface states at the ends of the fifth dimension, for $0 < m_0 < 2$ surface states for a single light quark flavor appear, for $2 < m_0 < 4$ surface states for four light flavors appear and for $4 < m_0 < 6$ there are surface states for six light quark flavors. There is a symmetry under $m_0 \rightarrow 10 - m_0$, which remains when coupling to gauge fields is introduced, although then these values of m_0 shift.

Figure 1 shows one way to probe the number of light fermions. The graph is for quenched simulations at $\beta = 5.85$ on an $8^3 \times 32$ lattice. For a given value of m_0 , we have measured the chiral condensate for $m_f = 0.02, 0.04, 0.06, 0.08$ and 0.10 , and calculated $\langle \bar{\psi}\psi(m_f \rightarrow 0) \rangle$ (simple linear fits work well). This extrapolation was done for three different values of L_s , to make sure we were close to the large L_s limit. The figure shows this extrapolated chiral condensate as a function of m_0 .

For $m_0 < 0.9$ the extrapolated chiral condensate is zero, indicating no spontaneous chiral symmetry breaking, due to the absence of surface states. For larger values of m_0 the chiral condensate is non-zero and increasing, until leveling off for $1.65 < m_0 < 2.15$. This gives a region where there appears to be a single surface state. (For free fermions, the normalization of the fields depends on m_0 ; this region of insensitivity to m_0 does not appear to have been anticipated.) For

*R. Mawhinney presented talk at Lattice '98. Work supported in part by the US Department of Energy. Calculations were done on the QCDSF computer at Columbia University.

†Supported by the Max Kade Foundation

‡Current address: Physics Department, University of Illinois, Urbana, IL 61801




Figure 1. The four-dimensional chiral condensate extrapolated to zero quark mass as a function of the five-dimensional domain wall mass.

larger m_0 the chiral condensate increases as the change from a one flavor to a four flavor theory occurs. Above about 3.4, a region of relative insensitivity to m_0 occurs, where there are four flavors of light fermions. The chiral condensate in this region is numerically very close to four times its value for $1.65 < m_0 < 2.15$.

A second basic question is whether the mixing between the chiral modes at $s = 0$ and $s = L_s - 1$ can be controlled by choosing L_s large. It is known that a gauge field with a zero eigenvalue for the four-dimensional Wilson Dirac operator (with mass m_0), has no damping for modes propagating in the fifth dimension. We wanted to explicitly measure the effect of this undamped mode on non-anomalous low energy QCD physics. To this end, we have studied its effect on the chiral condensate (for related work see [6]).

The four-dimensional chiral condensate is defined through the inverse propagator for domain wall fermions, where $s = 0(L_s - 1)$ is a source for right (left) handed quarks and $s = L_s - 1(0)$ is a sink for left (right) handed quarks. We have studied a generalized condensate, where the distance from source to sink is variable. When this distance is $L_s - 1$, we have the conventional condensate.

For a few thermalized configurations, we have measured the values of m_0 where there should be undamped modes in the fifth dimension. Figure 2 shows a measurement of the generalized chiral condensate as a function of the separation between the source and sink in the fifth dimension for m_0 very close to the value where a zero of the four-dimensional Wilson operator occurs.




Figure 2. A generalized version of the four-dimensional chiral condensate, where the separation in the fifth dimension between the source and sink is s . $s = L_s - 1$ is the usual condensate.

One can clearly see the presence of the expected translationally invariant mode in the fifth dimension. However, this mode contributes about 1% to the value of the chiral condensate, for this one lattice where its effect should be largest. For an ensemble of lattices, at any given m_0 only some will have largely undamped modes. (This number could be a set of measure zero in the full average, although this needs to be studied as a function of lattice volume.) Even if undamped modes do survive the ensemble average, our example shows

that they can make a very small contribution to an observable.

3. HADRON MASSES

To test the formulation in a more practical way, we have measured hadron masses for quenched domain wall fermions on $8^3 \times 32$ lattices with $L_s = 10, 16, 24$ and 48 for $\beta = 5.7$ and $m_0 = 1.65$. (Since the conference, we have also simulated at $L_s = 32$.) We have chosen m_f in the range 0.02 to 0.22 . For $m_f = 0.02$, we have not gotten good plateaus with 70-90 configurations and consequently only report results using $m_f > 0.02$.

The rho mass shows no dependence on L_s between 10 and 48, while the nucleon mass decreases somewhat for larger L_s . For $L_s = 10$, we find $m_\rho = 0.787(12) + 2.28(5)m_f$ and $m_N = 1.19(3) + 3.71(15)m_f$. For $L_s = 48$, we find $m_\rho = 0.792(28) + 2.15(12)m_f$ and $m_N = 1.12(4) + 3.79(19)m_f$. These are correlated fits to five or more valence quark masses and all have $\chi^2/dof < 1$. Figure 3 shows the nucleon and rho masses for $L_s = 48$. The $m_f = 0$ value for m_N/m_ρ is 1.42 ($L_s = 48$), compared to the staggered fermion values of ≈ 1.50 [7].




Figure 3. Rho ($\square$) and nucleon ($\diamond$) masses versus m_f for 80 quenched $8^3 \times 32$ configurations at $\beta = 5.7$. The fifth dimension has $L_s = 48$.

We have also measured $m_\pi(m_f)$ for these lattices and find that it is not a simple linear function of m_f for the full range of quark masses we used. Fits including an m_f^2 term, or linear fits to the few smallest values for m_f , give values for $m_\pi^2(m_f \rightarrow 0)$ that agree within errors. We then find that $m_\pi^2(0)$ fits well to the form $A \exp(-\alpha L_s) + B$, with $B \approx 0.05$.

We have not yet established an explanation for this non-zero intercept for m_π^2 . Our studies of the condensate make residual mixing in the fifth dimension seem unlikely. Possible causes include the finite volume effects which give the pion a non-zero intercept in quenched staggered spectroscopy, a problem with quenching that is brought out by the zero modes of domain wall fermions or the observed failure of our integrated π - π correlator to equal $\langle \bar{\psi}\psi \rangle / m_f$ for small values of m_f .

4. CONCLUSIONS

Our studies of domain wall fermions for quenched QCD have shown the technique capable of reproducing expected physics. The non-zero value for $m_\pi^2(0)$, even for large L_s , deserves more attention. Since other formulations have shown similar effects for $m_\pi^2(0)$, we must check again at weaker coupling and larger volumes before concluding that this is a new phenomena revealed by domain wall fermions in the quenched approximation.

The authors would like to thank Robert Edwards, Tim Klassen, Rajamani Narayanan and Tom Blum for useful discussions.

REFERENCES

1. D. Kaplan, Phys. Lett. **B288** (1992) 342.
2. V. Furman and Y. Shamir, Nucl. Phys. **B439** (1995) 54.
3. P. Vranas, Phys. Rev. **D57** (1998) 1415.
4. T. Blum and A. Soni, Phys. Rev. **D56** (1997) 174; Phys. Rev. Lett. **79** (1997) 3595.
5. T. Blum, these proceedings.
6. R. Edwards, *et. al.* hep-lat/9802016.
7. C. Bernard, *et. al.* Nucl. Phys. **B60A** (Proc. Suppl.) (1998) 3.